

Non-Invasive Imaging of Pediatric Abdominal Tumors: The Role of Ultrasound and Complementary Modalities

(Authors Details)

Sayed Ali¹, Samira Khan^{2*}, Imtiaz Ali³

¹MBBS, Pediatric, Resident (Pakistan) Department of Radiology, ACE Institute of Technology, Elmhurst, Queens, NY, USA

²RDMS, Lead Instructor Sonography ACE Institute of Technology, Elmhurst, Queens, NY, USA

³MD, DMRD, ARDMS, FACS, Professor of Diagnostic Medical Imaging, ACE Institute of Technology, Elmhurst, Queens, NY, USA

Email: drsamira@yahoo.com

Abstract

Background: Pediatric abdominal tumors present a diagnostic challenge due to nonspecific clinical symptoms and the need to minimize invasive procedures and exposure to ionizing radiation. Imaging is central to early detection, tumor characterization, and clinical management.

Objective: To review the role of non-invasive imaging in pediatric abdominal tumors, with a focus on ultrasound as a first-line diagnostic modality, and to compare its utility with other non-invasive techniques, including magnetic resonance imaging and computed tomography.

Methods: A narrative review of peer-reviewed literature was conducted, emphasizing studies on diagnostic accuracy, safety, and clinical applications of non-invasive imaging in pediatric abdominal tumors. Imaging guidelines, consensus statements, and recent technological advances were included

Results: Ultrasound is a highly effective, safe, and accessible first-line imaging tool for initial evaluation. Its real-time imaging, absence of ionizing radiation, and high diagnostic accuracy make it valuable for early detection and follow-up. Magnetic resonance imaging provides superior soft-tissue characterization and staging, whereas computed tomography is reserved for selected cases despite radiation concerns.

Conclusion: Non-invasive imaging is essential for diagnosing and managing pediatric abdominal tumors. Ultrasound, complemented by magnetic resonance imaging and computed tomography when indicated, provides a child-safe strategy that improves diagnostic confidence, guides clinical decision-making, and enhances patient outcomes. Advances in technology and standardized imaging protocols are expected to further optimize pediatric tumor evaluation.

Keywords: Pediatric abdominal tumors; non-invasive imaging; diagnostic ultrasound; pediatric oncology; abdominal imaging; magnetic resonance imaging; computed tomography

DOI: 10.46682/ijhbs.9.1.01

1. Introduction

Pediatric abdominal tumors represent a heterogeneous group of neoplastic conditions affecting organs such as the kidneys, liver, adrenal glands, gastrointestinal tract, ovaries, and retroperitoneum. These tumors often present with nonspecific symptoms, including abdominal distension, pain, or palpable masses, making early diagnosis challenging. Timely and accurate detection is critical, as delays may negatively affect treatment and prognosis.

Imaging is central to pediatric oncology, enabling tumor detection, localization, and characterization while guiding management. Children are particularly vulnerable to ionizing radiation due to increased tissue radiosensitivity and longer life expectancy, making non-invasive, radiation-free modalities the preferred choice.

Ultrasound is widely accepted as the first-line imaging tool because of its safety, accessibility, portability, and real-time capability. Advances in ultrasound and magnetic resonance imaging have enhanced tumor characterization, staging, and treatment monitoring. This review highlights the principles, applications, and limitations of non-invasive imaging in pediatric abdominal tumors, focusing on ultrasound-based strategies.

2. Pediatric Abdominal Tumors: Clinical Overview

2.1 Classification and Common Tumor Types

Pediatric abdominal tumors can be benign or malignant, with malignancies occurring more frequently in early childhood. Common malignant tumors include neuroblastoma, Wilms tumor, hepatoblastoma, lymphoma, and soft-tissue sarcomas, while benign lesions include ovarian cysts, mesenteric cysts, and hepatic masses.

Tumor origin strongly influences imaging appearance and clinical behavior. Wilms tumor usually presents as a well-circumscribed renal mass, whereas neuroblastoma often arises from the adrenal gland and may show calcifications and infiltrative growth. Ovarian masses in children range from functional cysts to neoplastic lesions, requiring careful imaging evaluation to guide management.

2.2 Epidemiology and Age-Related Presentation

The incidence and presentation of pediatric abdominal tumors vary according to age and tumor type. Neuroblastoma is the most common extracranial solid tumor in children under five years, while Wilms tumor predominantly affects those between two and five years. Hepatoblastoma primarily occurs in infants and young children. Younger children often present with rapidly enlarging abdominal masses, whereas older children may exhibit nonspecific symptoms such as abdominal pain, weight loss, or gastrointestinal disturbances. Age-related anatomical and

physiological differences must be considered when selecting imaging modalities and interpreting findings.

2.3 Diagnostic Challenges

The nonspecific clinical presentation of pediatric abdominal tumors frequently leads to delayed diagnosis. Imaging is therefore essential for early detection and differentiation between benign and malignant lesions. Challenges include patient cooperation, operator dependence, and limitations of individual imaging modalities. These factors underscore the importance of structured, multimodal imaging pathways tailored to pediatric patients.

3. Non-Invasive Imaging Principles in Pediatric Oncology

3.1 Definition and Clinical Advantages of Non-Invasive Imaging

Non-invasive imaging allows visualization of anatomy and pathological processes without penetrating the skin or inserting equipment into the body. It is fundamental for tumor diagnosis, characterization, staging, and follow-up in pediatric oncology, providing clinically useful information while minimizing physical harm. Common non-invasive modalities include ultrasound (US), magnetic resonance imaging (MRI), and computed tomography (CT), with US and MRI preferred due to their superior safety profiles [4–6].

These modalities enable early tumor detection, assessment of size and location, evaluation of internal structure, and identification of complications such as hemorrhage or organ displacement. US offers real-time imaging, portability, and accessibility, making it the first-line diagnostic tool, particularly for abdominal and duodenal evaluation, whereas MRI provides high-resolution soft-tissue imaging without ionizing radiation exposure [6,11]. Non-invasive imaging also facilitates longitudinal monitoring, allowing clinicians to assess tumor regression, recurrence, or complications over time. Long-term follow-up is often required in pediatric oncology, making the avoidance of cumulative diagnostic risks a key consideration [3,14].

3.2 Safety Considerations in Pediatric Populations

Safety is a core concern in pediatric imaging, as children are more sensitive to the effects of ionizing radiation and have a higher lifetime risk of radiation-induced malignancies. This has led to the prioritization of modalities that minimize or avoid radiation exposure [5,6,12]. US is recognized as the safest imaging modality due to its non-ionizing nature and absence of harmful biological effects when properly used, making it highly suitable for initial assessment, follow-up, and screening of pediatric abdominal tumors.

MRI is also considered safe, though sedation may be required in younger children to ensure image quality, introducing additional considerations [11]. CT is diagnostically valuable in selected cases, particularly when urgent evaluation of complex anatomy or acute complications

is needed, but its use is carefully managed due to radiation exposure. Imaging guidelines emphasize dose optimization and appropriate modality selection to ensure patient safety [4,5].


3.3 Role of Imaging in Diagnostic Decision-Making

Non-invasive imaging plays a decisive role in diagnostic decision-making throughout the clinical course of pediatric abdominal tumors. Imaging often provides the first objective evidence guiding clinical suspicion, referral pathways, and further evaluation. The ability to differentiate benign from malignant lesions influences decisions regarding biopsy, surgical intervention, or conservative management [3,8].

Imaging also supports tumor staging and treatment planning by defining local extension, vascular involvement, lymph node enlargement, and distant metastases. Accurate staging is essential for selecting therapeutic strategies and predicting outcomes. Multimodal imaging allows integration of information from US, MRI, and CT to achieve comprehensive assessment [6,11].

Furthermore, imaging findings contribute to interdisciplinary decision-making within pediatric oncology teams. Radiologic assessment informs discussions among pediatric oncologists, surgeons, and radiotherapists, ensuring evidence-based, individualized management. The central role of non-invasive imaging in guiding diagnosis and management underscores its importance as a cornerstone of pediatric oncologic care [4,8,14].

Bar Chart: Advantages of Non-Invasive Imaging Modalities in Pediatric Oncology







Figure1,2,3 Description

Bar chart comparing key advantages of non-invasive imaging modalities in pediatric oncology, including ultrasound, magnetic resonance imaging, and computed tomography. Metrics displayed include safety profile, diagnostic accuracy, accessibility, and suitability for repeated follow-up. Ultrasound demonstrates the highest overall safety and accessibility scores, while magnetic resonance imaging shows superior soft-tissue characterization, and computed tomography provides rapid anatomical assessment with lower safety scores due to radiation exposure.

4. Ultrasound as a First-Line Imaging Modality

Ultrasound (US) is widely adopted as the first-line imaging modality for evaluating pediatric abdominal tumors due to its safety, accessibility, and diagnostic versatility. US uses high-frequency sound waves to generate real-time images of internal structures without exposing patients to ionizing radiation. This feature is particularly important in pediatric oncology, allowing repeated examinations while minimizing long-term radiation risks [1,2,7]. Advances in US technology, including high-resolution transducers, Doppler techniques, and enhanced image-processing algorithms, have improved tissue differentiation and vascular assessment, thereby increasing diagnostic confidence [1,2,7].

High-resolution US enables detailed visualization of abdominal organs and masses, allowing assessment of tumor size, location, margins, internal composition, and relationships to adjacent structures. Doppler US contributes by evaluating tumor vascularity, which can aid in differentiating benign from malignant lesions and in estimating tumor aggressiveness. These features make US highly effective for initial tumor detection and for guiding decisions on further cross-sectional imaging or invasive procedures in children presenting with abdominal symptoms or palpable masses [3,5].

The diagnostic accuracy of US in pediatric abdominal tumors is well documented. Studies report high sensitivity in detecting masses in solid organs such as the kidneys, liver, and adrenal glands. US can identify characteristic features of common pediatric tumors, including the well-defined margins of Wilms tumor, the heterogeneous appearance of neuroblastoma, and the cystic or solid components of ovarian masses [1,9,13]. In many clinical settings, US serves as the primary imaging tool that determines the need for additional imaging or biopsy.

Despite its advantages, US has limitations. Image quality and diagnostic accuracy are operator-dependent and may be affected by patient cooperation, body habitus, and bowel gas. Deep-seated lesions or tumors in anatomically complex regions may be incompletely visualized, and US provides limited capability for comprehensive tumor staging or detection of distant metastases. In such cases, complementary modalities such as magnetic resonance imaging (MRI) or computed tomography (CT) are necessary to complete diagnostic assessment and staging [5,6,11].

US plays a critical role in the early detection and characterization of pediatric abdominal tumors. Early diagnosis facilitated by US supports timely referral and initiation of treatment, which are key determinants of favorable clinical outcomes. It is also widely used in treatment monitoring and follow-up, allowing assessment of tumor response and detection of recurrence without repeated radiation exposure [2,4,14]. Overall, US remains an essential component of non-invasive imaging in pediatric oncology and, when integrated into a multimodal strategy, significantly contributes to accurate diagnosis, effective patient management, and improved clinical outcomes in children.

5. Complementary Non-Invasive Imaging Modalities

Magnetic resonance imaging (MRI) plays a complementary role in the evaluation of pediatric abdominal tumors, particularly when ultrasound findings are inconclusive or detailed anatomical characterization is required. MRI provides superior soft-tissue contrast, enabling precise delineation of tumor borders, internal structure, and relationships to adjacent organs. This information is critical for tumor staging, surgical planning, and treatment response assessment [5,6,11]. MRI is especially valuable for complex hepatic, retroperitoneal, pelvic, and gastrointestinal tumors, where multiplanar imaging can increase diagnostic accuracy. The absence of ionizing radiation allows repeated examinations during long-term follow-up, making MRI a safe modality for children. Advanced techniques such as diffusion-weighted imaging and contrast-enhanced sequences further improve lesion characterization by providing information on tissue cellularity and vascularity, supplementing structural assessment from ultrasound [5,6].

However, MRI has limitations. Imaging sessions are lengthy and may require sedation in younger children to maintain image quality, introducing logistical and safety considerations. Limited availability and higher cost compared to ultrasound also restrict its use as a first-line modality [6,12].


Computed tomography (CT) remains a useful diagnostic tool in selected cases, particularly for emergency assessment, evaluation of tumor-related complications, or when MRI is unavailable or contraindicated. CT provides high spatial resolution, rapid image acquisition, and excellent visualization of calcifications, vascular anatomy, and acute complications such as hemorrhage or bowel obstruction [3,11]. Despite these advantages, CT involves ionizing radiation, which poses a significant risk in pediatric patients due to increased radiosensitivity and lifetime malignancy risk. Therefore, CT use in pediatric oncology follows principles of dose optimization, with child-specific protocols designed to minimize radiation while maintaining diagnostic quality [4,5].

The relative strengths and limitations of ultrasound, MRI, and CT underscore the importance of a multimodal imaging approach. Ultrasound serves as a safe, accessible first-line modality for detection and follow-up. MRI provides detailed soft-tissue characterization and accurate staging without radiation exposure, whereas CT offers rapid, high-resolution imaging for calcifications


and acute pathologies. Optimal diagnostic practice depends on integrating imaging findings with clinical history and patient-specific factors [4,6,8,14]. The complementary use of these non-invasive modalities enhances diagnostic accuracy, informs treatment planning, and improves clinical outcomes in pediatric oncology.

Box-and-Whisker Plots: Comparative Performance of Imaging Modalities in Pediatric Abdominal Tumors

a



b






Figure 4,5 Description:

Box-and-whisker plots presenting a qualitative comparison of non-invasive imaging modalities used in the evaluation of pediatric abdominal tumors. The figure summarizes relative trends in diagnostic performance for ultrasound and magnetic resonance imaging based on reported diagnostic accuracy, soft-tissue characterization capability, and suitability for repeated follow-up. Ultrasound demonstrates consistent performance with a narrow interquartile range, reflecting high safety and accessibility, while magnetic resonance imaging shows higher median values for soft-tissue characterization, supporting its complementary role in detailed tumor assessment.

6 Clinical Applications, Outcomes, and Limitations

Non-invasive imaging modalities are central to the clinical management of pediatric abdominal tumors, playing a critical role in diagnosis, staging, treatment planning, and follow-up. Ultrasound is the preferred first-line imaging tool due to its safety, accessibility, and ability to detect abdominal masses. Findings from ultrasound guide further decision-making regarding the need for cross-sectional imaging, biopsy, or referral to specialized oncology services [5,11,14].

Magnetic resonance imaging (MRI) provides additional anatomical information when needed, allowing precise assessment of tumor extent, local invasion, and regional lymph node involvement. Computed tomography (CT) is reserved for select cases, such as emergency presentations, evaluation of calcifications, or acute complications. Imaging directly informs treatment planning, enabling assessment of surgical feasibility, determination of neoadjuvant

therapy requirements, and risk stratification. Stepwise visualization of tumor boundaries, vascular involvement, and organ displacement aids surgeons and oncologists in selecting optimal therapeutic strategies while minimizing procedural risks.

Non-invasive imaging is equally critical for monitoring treatment response, facilitating evaluation of tumor regression, residual disease, or recurrence without subjecting children to cumulative radiation exposure. Radiation-free modalities like ultrasound and MRI enhance patient safety by reducing long-term health risks, including secondary malignancies. Furthermore, non-invasive imaging minimizes the need for invasive diagnostic procedures, decreasing procedure-related complications, hospitalization, and psychological trauma for pediatric patients and their families [5,11,14].

Despite these advantages, limitations exist. Ultrasound is operator-dependent, and image quality may be affected by examiner experience, patient cooperation, and technical factors such as body habitus or bowel gas. MRI, while highly informative, is limited by cost, availability, and the potential need for sedation in younger children. CT, although diagnostically valuable, carries radiation hazards that restrict its routine use in pediatric oncology. These considerations underscore the importance of standardized imaging guidelines, multidisciplinary collaboration, and careful selection of imaging modalities to optimize diagnostic accuracy and patient care [4,6,8].

In summary, integrating non-invasive imaging into the diagnostic and management pathway is essential for achieving accurate diagnosis, guiding treatment, and ensuring patient safety in pediatric abdominal tumors.

Conclusion

Non-invasive imaging has become a cornerstone in the diagnosis and management of pediatric abdominal tumors, providing a balance between diagnostic accuracy and patient safety. It plays a critical role in addressing the unique challenges of pediatric oncology, where early detection, precise tumor characterization, and careful staging are essential for optimal clinical outcomes. The integration of non-invasive modalities into diagnostic pathways has significantly enhanced the evaluation of abdominal tumors while minimizing procedural risks and exposure to ionizing radiation.

Ultrasound remains the first-line imaging modality due to its accessibility, real-time imaging capability, and excellent safety profile. Its utility in initial tumor detection and longitudinal follow-up makes it indispensable in both resource-rich and resource-limited settings. Magnetic resonance imaging (MRI) serves as a valuable complementary tool, offering superior soft-tissue contrast and detailed anatomical assessment to support accurate staging and treatment planning. Computed tomography (CT) is reserved for select clinical scenarios requiring rapid assessment

or evaluation of calcifications and acute complications, with careful attention to minimizing radiation exposure.

The judicious use of non-invasive imaging directly influences clinical decision-making, treatment planning, and long-term follow-up. Early and accurate imaging-guided diagnosis facilitates timely intervention, reduces reliance on invasive procedures, and contributes to improved survival and quality of life in pediatric patients. Awareness of modality-specific limitations, including operator dependence, sedation requirements, and logistical constraints, emphasizes the importance of standardized protocols and multidisciplinary collaboration.

In conclusion, a tailored, multimodal, non-invasive imaging strategy is essential for the effective evaluation of pediatric abdominal tumors. Ongoing technological advancements, coupled with evidence-based imaging guidelines, are expected to further enhance diagnostic precision, patient safety, and overall outcomes in pediatric oncology.

REFERENCES

1. Adedayo AA, Igashi JB, Mshelbwala PM, Nasir AA, Ameh EA, Adeniran JO. Accuracy of ultrasonography in the evaluation of abdominal masses in children in Nigeria. *Afr J Paediatr Surg.* 2019;16(1-4):1-5.
2. Handique M, Gogoi BJ, Biswas SS. Ultrasonographic evaluation of pediatric abdominal masses. *Asian J Med Sci.* 2025;16(9):143-147.
3. Garg HK, Shashi KK, Fisher P, Winant AJ, Hull NC, Lee EY. Pediatric upper abdominal masses: current practical imaging assessment. In: *Seminars in Roentgenology.* WB Saunders; 2024 Jul. Vol. 59(3):299-311.
4. Smriti V, Baheti AD, Shah S, Qureshi SS, Shetty N, Gala K, Kapadia T, et al. Imaging recommendations for diagnosis, staging, and management of pediatric solid tumors. *Indian J Med Paediatr Oncol.* 2023;44(1):138-148.
5. Kim HH, Hull NC, Lee EY, Phillips GS. Pediatric abdominal masses: imaging guidelines and recommendations. *Radiol Clin.* 2022;60(1):113-129.
6. Morin CE, Artunduaga M, Schooler GR, Brennan RC, Khanna G. Imaging for staging of pediatric abdominal tumors: an update, from the AJR special series on cancer staging. *AJR Am J Roentgenol.* 2021;217(4):786-799.
7. Stein NR, VanHouwelingen L. Making good use of ultrasound for abdominal tumors in children. *J Pediatr.* 2023;99(1):1-3.

8. Srinivasan A, Parikh A, Pace E, Schechter A, Tang E, Servaes S. Imaging of pediatric abdominal soft tissue tumors: a COG Diagnostic Imaging Committee/SPR Oncology Committee white paper. *Pediatr Blood Cancer*. 2023;70:e30341.
9. Özcan HN, Yavuz ÖÖ, Ekinci S, Oguz B, Kutluk T, Haliloglu M. Imaging findings of gastrointestinal tract tumors in children and adolescents. *Insights Imaging*. 2022;13(1):51.
10. Seidel FG. Imaging of a pediatric abdominal mass. In: *Seminars in Surgical Oncology*. New York: John Wiley & Sons; 1986. Vol. 2(3):125-138.
11. Jha SK, Brown C, Kang L, Diaz ES, Gwal K, Alvarez E, Stein-Wexler R, et al. Update on the role of imaging in staging of common pediatric abdominal tumors. *Curr Probl Cancer*. 2023;47(2):100969.
12. Clayton KM. Focus on diagnosis: pediatric abdominal imaging. *Pediatrics Rev*. 2010;31(12):506-510.
13. Bourgioti C, Konidari M, Giantsouli A, Kafritsa A, Xydis V, Mouloupoulos LA, Tsili AC, et al. Imaging evaluation of ovarian masses in a pediatric population: a comprehensive overview. *Cancers*. 2025;17(14):2316.
14. Davis LM, Martinez-Correa S, Freeman CW, Adams C, Sultan LR, Le DQ, Hwang M, et al. Ultrasound innovations in abdominal radiology: techniques and clinical applications in pediatric imaging. *Abdom Radiol*. 2025;50(4):1744-1762.
15. Bahinipati P, Sahoo RK, Mohanty BB, Das SR, Panda SK. Computed tomographic evaluation of paediatric abdominal mass lesions. *Int J Pharm Sci Rev Res*. 2016;37(1):175-179.