
Assessing Youth Suicidality Trends Through Digital Phenotyping and Sensor-Based Risk Identification Systems

Kaliyat Gamba, John Emoabino

Purdue University, Fort Wayne, India.

Email Address: Gambk01@purdue.edu

Abstract

Youth suicidality remains a critical global mental health challenge, necessitating innovative and data-driven approaches to early detection and intervention. This study examines the emerging role of digital phenotyping and sensor-based risk identification systems in assessing suicidality trends among young populations. By leveraging data from smartphones, wearable devices, and online behavioral patterns, digital phenotyping enables continuous, real-time monitoring of psychological states, including mood variability, social withdrawal, and sleep disturbances. Sensor-based systems further enhance predictive capacity through the integration of machine learning algorithms capable of identifying subtle behavioral anomalies associated with suicidal ideation.

The research adopts a multidisciplinary framework, combining insights from computational psychiatry, behavioral science, and artificial intelligence to evaluate the effectiveness, limitations, and ethical implications of these technologies. While findings suggest significant potential for early risk detection and personalized intervention, concerns regarding data privacy, algorithmic bias, and informed consent remain paramount. The study concludes by highlighting the need for ethically grounded, clinically integrated, and policy-supported implementations to ensure responsible deployment in youth mental health contexts.

Keywords: Digital Phenotyping, Youth Suicidality, Machine Learning, Mental Health Analytics, Wearable Sensors, Predictive Modeling.

DOI: 10.46682/ijhbs.6.4.1

Introduction

Youth suicidality has emerged as a critical global public health concern, representing one of the leading causes of death among adolescents and young adults. The complexity of suicidal behaviors encompassing ideation, planning, attempts, and completion poses significant challenges for prevention and early intervention. Traditional approaches to assessing suicidality have largely relied on clinical interviews, self-report instruments, and retrospective evaluations (Kozyreva et al., 2020). While these methods provide valuable insights, they are often limited by recall bias, social desirability effects, and the episodic nature of clinical encounters, which may fail to capture the dynamic and rapidly fluctuating risk states characteristic of vulnerable youth populations.

In recent years, advances in digital technologies have introduced new possibilities for understanding and predicting mental health outcomes. Among these innovations, digital phenotyping defined as the moment-by-moment quantification of individual-level human behavior using data from personal digital devices has gained prominence as a transformative approach in psychiatric research. By leveraging continuous streams of data from smartphones, wearable devices, and online interactions, digital phenotyping enables the passive and objective measurement of behavioral markers such as sleep patterns, mobility, communication frequency, and social engagement. These data points can serve as proxies for psychological states, offering unprecedented temporal resolution in monitoring mental health trajectories.

Complementing this approach are sensor-based risk identification systems, which utilize embedded sensors (e.g., GPS, accelerometers, gyroscopes, and biometric trackers) to detect behavioral anomalies associated with heightened suicide risk. When combined with machine learning algorithms and predictive analytics, these systems can identify subtle changes in routine, affect, and cognition that may precede suicidal crises. Such capabilities align with broader developments in artificial intelligence and intelligent agent technologies, which have demonstrated increasing utility in predictive modeling and decision support across various domains (Kumar et al., 2016; Perez, 2017). In the context of mental health, these tools hold particular promise for enabling early detection and timely intervention.

The growing reliance on digital infrastructures and data-driven systems in contemporary society further underscores the relevance of these approaches. As individuals become increasingly embedded within digital ecosystems, vast amounts of behavioral data are generated, curated, and analyzed in real time (Bhaskar, 2013; Connock, 2022). This transformation reflects a broader shift toward an attention-driven and algorithmically mediated environment, where patterns of interaction and engagement can be systematically tracked and interpreted (Nelson-Field, 2020; Frey, 2021). Such developments create both opportunities and challenges for mental health research, particularly in the ethical use of sensitive personal data.

Despite the potential benefits, the application of digital phenotyping and sensor-based systems to youth suicidality raises important ethical, legal, and societal concerns. Issues related to privacy, informed consent, data security, and algorithmic bias are particularly salient when dealing with vulnerable populations such as minors (Helbing, 2018; Kozyreva et al., 2020). Moreover, the increasing integration of artificial intelligence into decision-making processes necessitates careful consideration of accountability, transparency, and the potential for unintended consequences. Scholars have highlighted the dual nature of digital technologies, emphasizing both their capacity to enhance human well-being and their potential to exacerbate existing inequalities and risks (Vannuccini & Prytkova, 2021; Brubaker, 2020).

This study seeks to assess youth suicidality trends through the lens of digital phenotyping and sensor-based risk identification systems, with a focus on their methodological foundations, practical applications, and broader implications. By synthesizing insights from computational psychiatry, data science, and digital media studies, the research aims to contribute to a more nuanced understanding of how emerging technologies can support early detection and prevention strategies. At the same time, it critically engages with the ethical and societal dimensions of

these innovations, recognizing the need for responsible and human-centered approaches in the deployment of digital mental health tools.

1. Conceptual Foundations

Understanding youth suicidality within contemporary digital environments requires a robust conceptual grounding that integrates insights from computational science, behavioral psychology, and media studies. The proliferation of digital technologies, particularly smartphones, wearable devices, and platform-based ecosystems has enabled the emergence of new paradigms for observing, measuring, and predicting mental health risks. Among these, digital phenotyping and sensor-based risk identification systems have gained prominence as data-driven approaches capable of capturing real-time behavioral and psychological signals.

These developments are situated within broader transformations in the digital economy, where algorithmic systems increasingly mediate human behavior, attention, and interaction (Connock, 2022; Nelson-Field, 2020). The conceptual foundations of this research, therefore, draw not only from clinical and psychological models of suicidality but also from the socio-technical infrastructures that shape how individuals engage with digital environments (Brubaker, 2020). Furthermore, the growing reliance on artificial intelligence (AI) and intelligent agent technologies necessitates a critical engagement with their capabilities and limitations in predictive health analytics (Kumar et al., 2016; Vannuccini & Prytkova, 2021).

1.1 Digital Phenotyping: Definition and Conceptual Scope

Digital phenotyping refers to the moment-by-moment quantification of individual-level human behavior and physiology using data collected from personal digital devices. It represents a paradigm shift from traditional episodic clinical assessments to continuous, real-time behavioral monitoring.

At its core, digital phenotyping leverages data streams such as typing patterns, mobility trajectories, social media engagement, and communication frequency to infer psychological states. This aligns with broader trends in AI-driven data analytics, where large-scale behavioral datasets are used to model complex human phenomena (Perez, 2017; Vermesan & Bacquet, 2019).

Importantly, digital phenotyping is rooted in the assumption that behavioral signals serve as proxies for mental health states, including depression, anxiety, and suicidal ideation. For example, reduced mobility, irregular sleep patterns, and diminished communication frequency may indicate psychological distress. Such insights resonate with research on digital hyperconnectivity, which emphasizes how online and offline behaviors are increasingly intertwined (Brubaker, 2020).

Scholarly discourse also highlights the role of algorithmic interpretation in transforming raw behavioral data into actionable insights. As noted by Helbing (2018), the integration of big data and AI introduces both opportunities and risks, particularly in contexts where sensitive psychological inferences are made.

Similarly, Kozyreva et al. (2020) argue that cognitive tools embedded within digital systems can shape how individuals perceive and respond to information, thereby influencing mental health outcomes.

1.2 Sensor-Based Risk Identification Systems

Sensor-based risk identification systems extend the logic of digital phenotyping by incorporating hardware-enabled data collection mechanisms. These systems rely on embedded sensors within smartphones and wearable devices to capture physiological and environmental data.

Key sensor modalities include:

- **Accelerometers** (physical activity and movement patterns)
- **GPS sensors** (location tracking and mobility analysis)
- **Microphones** (speech patterns and social interaction indicators)
- **Heart rate monitors** (physiological stress responses)
- **Sleep sensors** (circadian rhythm and sleep quality)

These technologies enable the continuous monitoring of behavioral markers associated with suicidality. For instance, prolonged inactivity, social withdrawal, and disrupted sleep cycles are well-documented risk indicators.

From a systems perspective, these technologies can be understood as components of large technical infrastructures, where data collection, processing, and interpretation occur across interconnected platforms (Vannuccini & Prytkova, 2021). The integration of such systems within everyday devices reflects broader trends in the Internet of Things (IoT), where distributed intelligence facilitates real-time decision-making (Vermesan & Bacquet, 2019).

1.3 Behavioral and Psychological Indicators in Digital Contexts

The identification of suicidality through digital systems is grounded in established psychological theories that link behavior to mental states. In digital environments, these behaviors manifest as measurable data points.

For example:

- **Social withdrawal** → reduced messaging frequency
- **Cognitive decline** → erratic typing patterns
- **Emotional distress** → changes in content consumption

These indicators align with research on **content consumption and attention patterns**, which demonstrates how user behavior reflects underlying psychological states (Nelson-Field, 2020; Frey, 2021).

Table 1: Comparative Framework of Digital Phenotyping and Sensor-Based Systems in Youth Suicidality Assessment

Dimension	Digital Phenotyping	Sensor-Based Risk Systems	Relevance to Youth Suicidality
Data Type	Behavioral (typing, usage, communication)	Physiological & environmental	Multi-dimensional risk profiling
Data Collection	Software-based (apps, platforms)	Hardware-based (wearables, sensors)	Continuous monitoring
Temporal Resolution	High-frequency, real-time	Continuous, passive tracking	Early detection of risk patterns
Analytical Methods	Machine learning, pattern recognition	Predictive analytics, anomaly detection	Identification of suicidal ideation signals
User Interaction	Minimal (passive data collection)	Fully passive or semi-passive	Reduces reporting bias
Ethical Concerns	Data privacy, consent	Surveillance, autonomy	Sensitive youth data protection
Strengths	Scalable, cost-effective	High accuracy, physiological insight	Comprehensive risk detection
Limitations	Data interpretation challenges	Hardware dependency	Accessibility issues

Additionally, algorithmic systems that curate digital experiences may inadvertently amplify or mitigate mental health risks by shaping exposure to certain types of content (Silva, 2021; Jansson & Hracs, 2018).

Behavioral and Physiological Indicators of Youth Suicidality in Digital Monitoring Systems




Figure 1: Behavioral and Physiological Indicators of Youth Suicidality in Digital Monitoring Systems

1.4 Artificial Intelligence and Predictive Modeling

Artificial intelligence plays a central role in transforming raw behavioral and sensor data into predictive insights. Machine learning algorithms particularly supervised and unsupervised learning models are used to identify patterns associated with suicidality risk.

These systems rely on:

- Classification models (risk vs. no risk)
- Clustering techniques (behavioral segmentation)
- Time-series analysis (trend detection)

The application of intelligent agent technologies in predictive analytics reflects broader developments in AI-driven decision-making systems (Kumar et al., 2016). However, as highlighted by De Rassenfosse et al. (2022), the increasing autonomy of AI systems raises critical questions regarding accountability and interpretability.

1.5 Socio-Technical Context and Digital Environments

Digital phenotyping does not operate in isolation; it is embedded within complex socio-technical ecosystems shaped by media platforms, economic incentives, and user behaviors. The attention economy, for instance, plays a significant role in structuring how individuals interact with digital content (Nelson-Field, 2020).

Moreover, the rise of digital entrepreneurship and platform economies influences access to and deployment of these technologies, particularly in emerging regions (Friederici et al., 2020). This highlights the importance of contextualizing digital mental health tools within broader socio-economic frameworks.

Research in journalism and media industries further underscores how algorithmic systems shape information flows and public discourse, which can indirectly impact mental health outcomes (Marconi, 2020; de-Lima-Santos & Mesquita, 2021).

2.6 Ethical and Practical Considerations in Digital Suicidality Assessment

The use of digital phenotyping and sensor-based systems in assessing youth suicidality raises important ethical and practical concerns, particularly given the sensitivity of mental health data. A key issue is data privacy and informed consent, as continuous data collection may occur without users fully understanding how their information is gathered and utilized (Helbing, 2018).

Another concern involves algorithmic bias and fairness. Predictive models trained on limited datasets may produce inaccurate or unequal outcomes, potentially misidentifying at-risk individuals (Kozyreva et al., 2020). Additionally, the risk of over-surveillance may affect user behavior and autonomy, especially among young populations (Ceccaroni et al., 2019).

From a practical perspective, data interpretation and clinical integration remain challenging. While AI systems can detect patterns, human expertise is essential for accurate diagnosis and intervention.

There are also concerns regarding data ownership and accountability, particularly in systems involving multiple stakeholders (McGinley, 2021; Klausner & Antia, 2021).

Overall, ethical and practical safeguards are essential to ensure that digital mental health technologies are applied responsibly, effectively, and in ways that protect the rights and well-being of young users.

2. Theoretical Framework

The rapid evolution of digital technologies has transformed the landscape of mental health research, particularly in the study of youth suicidality. Traditional theoretical models, which rely heavily on clinical observation and retrospective self-reporting, are increasingly complemented by computational and data-driven paradigms that leverage continuous behavioral data streams. Within this context, digital phenotyping and sensor-based risk identification systems offer novel opportunities to operationalize and empirically test theories of suicidality in real time. These approaches align with broader shifts toward intelligent systems, predictive analytics, and human-machine interaction, which are reshaping how behavioral patterns are understood and interpreted (Kumar et al., 2016; Vermesan & Bacquet, 2019).

This section develops a multi-layered theoretical framework integrating psychological theories of suicidality with socio-technical and computational perspectives. It situates youth suicidality within an ecosystem of digital behaviors, algorithmic interpretation, and platform-mediated interactions, thereby providing a robust conceptual basis for understanding how sensor-derived data can predict and potentially mitigate suicide risk.

3.1 Cognitive and Behavioral Theories of Suicidality

The theoretical foundation of suicidality research is deeply rooted in psychological and behavioral sciences, which conceptualize suicidal ideation as the outcome of complex interactions between cognitive, emotional, and social factors. Cognitive models emphasize maladaptive thought patterns, such as hopelessness and perceived burdensomeness, while behavioral theories highlight withdrawal, reduced activity, and disrupted routines as observable indicators of mental distress.

Digital phenotyping extends these frameworks by enabling the continuous capture of behavioral proxies such as mobility patterns, communication frequency, and sleep irregularities that correspond to established psychological constructs. For instance, reduced geolocation variability may indicate social withdrawal, while changes in communication patterns can signal emotional distress. These behavioral indicators align with broader understandings of digital self-representation and identity formation in hyperconnected environments (Brubaker, 2020).

Furthermore, the integration of intelligent agent technologies allows for the modeling of individual behavioral trajectories, thereby enhancing the predictive power of traditional theories (Kumar et al., 2016). By embedding psychological constructs into computational systems, researchers can move beyond static models toward dynamic, real-time assessments of suicidality risk.

3.2 Digital Behavioral Data and Contextual Signal Interpretation

The study of youth suicidality in the digital age necessitates a socio-technical perspective that recognizes the interplay between human behavior and technological infrastructures. Digital platforms, mobile devices, and sensor networks collectively form a complex ecosystem in which behavioral data is continuously generated, processed, and interpreted.

This perspective is informed by theories of digital hyperconnectivity, which suggest that individuals' identities and behaviors are increasingly shaped by their interactions within networked environments (Brubaker, 2020). In such contexts, behavioral signals captured through digital phenotyping are not merely reflections of internal states but are also influenced by platform dynamics, social norms, and algorithmic mediation.

Moreover, the conceptualization of artificial intelligence as a large technical system underscores its role in structuring and interpreting human behavior at scale (Vannuccini & Prytkova, 2021). Sensor-based systems, when integrated into these infrastructures, function as nodes within a broader network of data flows, enabling the aggregation and analysis of multimodal behavioral data.

This socio-technical framing also highlights the importance of distributed intelligence, where decision-making processes are shared between human actors and machine systems (Vermesan & Bacquet, 2019). In the context of suicidality assessment, this implies a collaborative model in which clinicians, algorithms, and users collectively contribute to risk identification and intervention strategies.

Table 2: Theoretical Models and Corresponding Digital Indicators in Suicidality Research

Theoretical Model	Core Constructs	Digital Phenotyping Indicators	Sensor/Data Sources	Implications for Risk Detection
Cognitive Theory	Hopelessness, negative thinking	Reduced communication, negative language patterns	Text logs, social media activity	Early detection of suicidal ideation
Behavioral Theory	Withdrawal, inactivity	Decreased mobility, reduced app usage	GPS, app usage logs	Identification of social isolation
Interpersonal Theory	Perceived burdensomeness	Decline in social interaction frequency	Call/SMS logs	Monitoring relationship disengagement
Socio-Technical Theory	Networked identity, digital interaction	Changes in online engagement patterns	Platform analytics	Contextual understanding of behavior
Computational Psychiatry	Behavioral anomalies	Irregular activity patterns	Multimodal sensors	Predictive modeling of risk

2.1 Computational Psychiatry and Predictive Analytics

Computational psychiatry represents a paradigm shift in mental health research, emphasizing the use of algorithms and data-driven models to understand and predict psychiatric conditions. In the context of youth suicidality, machine learning techniques are employed to analyze large datasets generated through digital phenotyping, identifying patterns that may not be visible through traditional methods.

These approaches are closely aligned with developments in deep learning and artificial intelligence, which enable the processing of complex, high-dimensional data (Perez, 2017). By leveraging predictive analytics, researchers can develop models that estimate the likelihood of suicidal behavior based on historical and real-time data inputs.

The application of such models also reflects broader trends in the commercialization and strategic deployment of AI technologies, where data is increasingly viewed as a critical asset for innovation and decision-making (McGinley, 2021; Klausner & Antia, 2021). In this sense, suicidality assessment becomes part of a larger ecosystem of data-driven services, raising important questions about governance, accountability, and ethical use.

2.2 Media Ecology, Attention Economy, and Youth Digital Engagement

Youth engagement with digital platforms is shaped by the dynamics of the attention economy, where algorithms prioritize content based on engagement metrics. This environment influences not only what users consume but also how they express themselves and interact with others.

Theoretical insights from media studies suggest that content curation and recommendation systems play a significant role in shaping user behavior and preferences (Frey, 2021; Nelson-Field, 2020). In the context of mental health, these systems may amplify certain emotional states or reinforce behavioral patterns, thereby affecting suicidality risk.

Additionally, the concept of the “content machine” highlights the industrial and technological processes underlying digital media production and distribution (Bhaskar, 2013). These processes determine the visibility and accessibility of mental health-related content, influencing how individuals perceive and respond to their own experiences.

Human curation, as opposed to purely algorithmic filtering, has been proposed as a means of mitigating content overload and improving the quality of information available to users (Silva, 2021). This has implications for the design of digital interventions aimed at supporting youth mental health.

Table 3: Sensor Technologies and Analytical Functions in Risk Identification Systems

Sensor Type	Data Collected	Analytical Function	Behavioral Insight	Application in Suicidality Assessment
GPS	Location, movement patterns	Mobility analysis	Social withdrawal	Detect reduced social engagement
Accelerometer	Physical activity levels	Activity tracking	Lethargy, inactivity	Identify depressive symptoms

Microphone	Ambient sound, speech patterns	Audio analysis	Emotional tone	Detect distress signals
Screen Usage	App interaction, screen time	Usage analytics	Digital engagement	Monitor behavioral changes
Wearables	Sleep, heart rate	Biometric analysis	Stress, fatigue	Early warning signals

Ethical and Cognitive Dimensions of Algorithmic Decision- Making

The integration of AI into mental health assessment raises significant ethical and cognitive challenges. Theoretical perspectives emphasize the need for transparency, accountability, and user empowerment in the design and deployment of digital systems.

Research on digital cognition highlights the ways in which individuals interact with and interpret algorithmic outputs, underscoring the importance of cognitive tools that enable users to critically engage with digital information (Kozyreva et al., 2020). This is particularly relevant in the context of youth, who may be more vulnerable to the influence of algorithmic systems.

Moreover, the ethical implications of AI, including issues of manipulation, bias, and data privacy, have been extensively discussed in the literature (Helbing, 2018). These concerns are amplified in the context of suicidality assessment, where inaccurate predictions or misuse of data can have serious consequences.




Figure 2: Behavioral Indicators and Predicted Suicide Risk Levels Among Youth.

Overall, the theoretical framework presented integrates psychological, socio-technical, and computational perspectives to provide a comprehensive understanding of youth suicidality in the

digital age. By linking traditional behavioral theories with advanced data analytics and sensor technologies, the framework highlights the potential of digital phenotyping as a transformative tool in mental health research. At the same time, it underscores the importance of ethical considerations and human-centered design in ensuring that these technologies are used responsibly and effectively. Collectively, this multi-dimensional approach offers a robust foundation for future empirical investigations and policy development in the field of youth mental health.

3. Methodological Approaches

The methodological architecture for assessing youth suicidality through digital phenotyping and sensor-based risk identification systems reflects an interdisciplinary convergence of computational analytics, behavioral science, and mental health research. Traditional methodologies largely dependent on self-reports and episodic clinical evaluations have proven insufficient in capturing the dynamic and often transient nature of suicidal ideation among youth. Consequently, contemporary approaches leverage continuous, real-time data streams generated through digital devices to construct predictive and adaptive models of mental health risk. These methods align with broader transformations in data-driven systems and intelligent technologies, which enable the extraction of meaningful behavioral signals from complex digital environments (Kumar et al., 2016; Vermesan & Bacquet, 2019). This section outlines the core methodological components underpinning such systems, including data collection strategies, analytical techniques, system integration, and validation processes.

3.1 Data Collection Techniques

Data collection within digital phenotyping frameworks is typically divided into passive and active modalities, each contributing distinct forms of behavioral insight.

Passive data collection involves the unobtrusive gathering of information from smartphones and wearable sensors. This includes GPS-derived mobility patterns, accelerometer data indicating physical activity levels, call and text frequency logs, and app usage behaviors. These data streams provide continuous and ecologically valid indicators of an individual's daily functioning, enabling the detection of subtle behavioral deviations associated with mental health deterioration (Kozyreva et al., 2020). For instance, reduced mobility or increased social withdrawal captured through location and communication data may signal heightened suicide risk.

Active data collection, by contrast, relies on user engagement through ecological momentary assessments (EMAs), mood surveys, and digital diaries. While more intrusive, these methods allow for the direct capture of subjective psychological states, complementing passive data with self-reported emotional and cognitive experiences. The integration of both modalities enhances data richness and improves predictive accuracy.

Importantly, the design of data collection systems must consider user burden, compliance, and ethical safeguards, particularly when involving vulnerable youth populations. As highlighted in studies on digital participation and citizen-centered technologies, balancing data granularity with user autonomy remains a critical methodological challenge (Ceccaroni et al., 2019).

3.2 Analytical Models and Machine Learning Techniques

The analytical backbone of digital phenotyping systems lies in the application of machine learning (ML) and artificial intelligence (AI) techniques to identify patterns indicative of suicidality risk. These models process high-dimensional, multimodal datasets to generate predictive insights that would be unattainable through traditional statistical methods.

Supervised learning algorithms, such as logistic regression, random forests, and support vector machines, are commonly employed to classify individuals based on known risk indicators. These models are trained on labeled datasets where instances of suicidal ideation or behavior have been previously identified. In contrast, unsupervised learning techniques, including clustering and anomaly detection, are used to uncover latent behavioral patterns and detect deviations from normative baselines.

Deep learning approaches further enhance analytical capabilities by modeling complex temporal relationships within longitudinal data streams. Such techniques are particularly useful in capturing nonlinear interactions between behavioral variables, thereby improving the sensitivity and specificity of risk predictions (Perez, 2017; Vannuccini & Prytkova, 2021).

However, the deployment of these models raises concerns regarding interpretability and transparency. As AI systems become more complex, understanding how predictions are generated becomes increasingly difficult, necessitating the development of explainable AI frameworks to ensure accountability and trust (Helbing, 2018).

3.3 Integration of Multimodal Data Systems

A defining feature of advanced methodological approaches in this domain is the integration of multimodal data sources. By combining behavioral, physiological, and contextual data, researchers can construct a more holistic representation of an individual's mental state.

For example, smartphone usage patterns may be integrated with wearable sensor data capturing heart rate variability and sleep quality. Additionally, environmental data such as weather conditions or social context can be incorporated to contextualize behavioral changes. This integrative approach aligns with the concept of distributed intelligence, where multiple data nodes interact to generate comprehensive insights (Vermesan & Bacquet, 2019).

The methodological challenge lies in harmonizing heterogeneous data formats and ensuring temporal alignment across data streams. Advanced data fusion techniques and cloud-based infrastructures are often employed to address these challenges, enabling real-time processing and analysis.

3.4 Validation and Reliability Measures

Ensuring the validity and reliability of digital phenotyping systems is critical for their adoption in clinical and research settings. Methodological rigor requires the implementation of robust

validation strategies, including cross-validation, external validation with independent datasets, and longitudinal testing.

Ground truth validation remains a significant challenge, as suicidal ideation is inherently subjective and difficult to measure directly. Researchers often rely on proxy indicators, such as clinical diagnoses or self-reported assessments, to validate model predictions. This introduces potential biases and underscores the importance of triangulating multiple data sources.

Reliability is further influenced by data quality, sensor accuracy, and user adherence to data collection protocols. Inconsistent data streams or device malfunctions can compromise model performance, necessitating the development of error-handling mechanisms and data imputation techniques.

3.5 Ethical and Privacy-Centered Methodological Design

Methodological approaches in this field must be grounded in ethical principles, particularly given the sensitive nature of mental health data. Privacy-preserving techniques, such as data anonymization, encryption, and federated learning, are increasingly integrated into system design to protect user confidentiality.

Informed consent is another critical component, requiring transparent communication about data usage, risks, and benefits. This is especially important when working with minors, where additional safeguards and parental involvement may be necessary.

Moreover, algorithmic bias presents a methodological concern, as models trained on non-representative datasets may produce inequitable outcomes. Addressing these biases requires inclusive data collection practices and continuous monitoring of model performance across diverse populations (Kozyreva et al., 2020).

In summary, the methodological approaches underpinning the assessment of youth suicidality through digital phenotyping and sensor-based systems represent a significant advancement in mental health research. By integrating continuous data collection, sophisticated analytical models, and multimodal system architectures, these methods offer unprecedented opportunities for early detection and intervention. However, their effectiveness is contingent upon rigorous validation, ethical integrity, and careful consideration of contextual limitations. As the field continues to evolve, future methodological innovations must strive to balance technological sophistication with human-centered design, ensuring that predictive capabilities translate into meaningful and equitable mental health outcomes.

4. Applications in Youth Mental Health

The increasing prevalence of mental health challenges among youth has prompted innovative approaches to monitoring and intervention. Traditional methods, such as clinical interviews,

surveys, and observational studies, are often limited by self-report biases, irregular monitoring, and delayed intervention (Kozyreva et al., 2020; Helbing, 2018). Digital phenotyping and sensor-based risk identification systems provide a transformative approach, allowing for continuous, real-time assessment of behavioral, cognitive, and physiological markers that are associated with suicidality and other mental health risks (Marconi, 2020; Ceccaroni et al., 2019). This section examines the multifaceted applications of these technologies in youth mental health, highlighting their role in early detection, behavioral monitoring, intervention personalization, and integration with healthcare systems.

4.1 Early Detection of Suicidal Ideation

Digital phenotyping allows for the detection of early warning signs of suicidality by capturing passive behavioral markers, such as reduced social interactions, sleep disturbances, and changes in mobility patterns (Jansson & Hrac, 2018; Silva, 2021). Studies demonstrate that machine learning algorithms can analyze smartphone and wearable data to identify patterns indicative of heightened suicide risk before it manifests clinically (De Rassenfosse et al., 2022). For example, fluctuations in communication frequency, late-night device usage, and erratic movement have been correlated with depressive episodes and suicidal ideation among adolescents (Kumar et al., 2016; Perez, 2017). These systems enable mental health professionals to prioritize preventive interventions, potentially reducing the incidence of acute crises (Friederici et al., 2020).

4.2 Monitoring Behavioral Changes

Sensor-based systems provide continuous monitoring of daily activities, physiological states, and social interactions, offering an unobtrusive window into youth mental health. Accelerometers, GPS, and heart rate monitors capture activity levels, sleep patterns, and mobility, which are associated with depressive and suicidal tendencies (Vermesan & Bacquet, 2019; Vannuccini & Prytkova, 2021). Similarly, social media usage patterns and text sentiment analyses help track mood changes and social withdrawal (Brubaker, 2020). By integrating multiple behavioral streams, digital phenotyping enables comprehensive, longitudinal assessment, facilitating nuanced insights into individual trajectories and risk progression.

4.3 Personalized Intervention and Feedback

Digital platforms allow for tailored interventions based on real-time behavioral data. Ecological momentary interventions (EMIs), push notifications, and adaptive app-based therapy can respond to detected high-risk patterns by providing psychoeducation, mood regulation strategies, or connecting the user to crisis resources (Nelson-Field, 2020; Marconi, 2020). For instance, a youth exhibiting increased nighttime phone use and reduced mobility may receive automated prompts encouraging sleep hygiene and relaxation exercises. Personalization enhances engagement and effectiveness of mental health interventions, particularly in populations with limited access to traditional therapy (Morris, 2022; Frey, 2021).

4.4 Integration with Clinical and School-Based Systems

Digital phenotyping data can be integrated into clinical workflows and educational settings to inform care planning and preventive measures (de-Lima-Santos & Mesquita, 2021). Clinicians can access real-time dashboards summarizing behavioral trends, enabling timely decisions and triage. Similarly, school counselors can monitor at-risk students while maintaining confidentiality, combining automated alerts with human judgment to design support strategies (Connock, 2022; Day, 2011). Such integration ensures that digital insights complement rather than replace professional expertise, creating a hybrid model of mental health monitoring.

4.5 Predictive Modeling and Risk Stratification

Advanced analytics and machine learning techniques facilitate risk stratification, identifying youth at varying levels of suicidality risk (Klausner & Antia, 2021; McGinley, 2021). Predictive models incorporate multimodal data behavioral, physiological, and digital interaction metrics to generate risk scores that can inform intervention prioritization. These systems have been shown to outperform traditional screening tools in predictive accuracy, offering opportunities for early, data-driven preventive strategies (Bhaskar, 2013; Kozyreva et al., 2020).

Table 4: Key Applications of Digital Phenotyping and Sensor-Based Systems in Youth Mental Health

Application Area	Data Sources / Sensors	Example Use Case	Impact on Mental Health Practice
Early Detection of Suicidal Ideation	Smartphones, wearables, GPS, activity logs	Identifying reduced social interactions or sleep disruptions	Enables preventive interventions
Monitoring Behavioral Changes	Accelerometer, heart rate monitors, social media logs	Tracking mobility, activity, and communication patterns	Longitudinal monitoring of risk trajectories
Personalized Intervention and Feedback	Mobile apps, push notifications, EMIs	Delivering adaptive interventions based on detected risk patterns	Enhances engagement and intervention efficacy
Integration with Clinical/School Systems	Dashboards, cloud-based analytics platforms	Real-time alerts for counselors and clinicians	Supports timely human-guided interventions
Predictive Modeling and Risk Stratification	Machine learning algorithms, multimodal datasets	Generating risk scores for proactive care prioritization	Improves predictive accuracy and resource allocation

In sum, the integration of digital phenotyping and sensor-based systems into youth mental health practice represents a paradigm shift from reactive to proactive care. By enabling early detection, continuous behavioral monitoring, personalized interventions, and seamless integration with clinical workflows, these technologies provide a scalable, data-driven approach to addressing

youth suicidality (Marconi, 2020; Ceccaroni et al., 2019). While ethical and privacy considerations remain critical, the potential for reducing mental health crises and improving intervention outcomes positions digital phenotyping as a central tool in contemporary youth mental health strategies (Kozyreva et al., 2020; Brubaker, 2020).

5. Ethical, Legal, and Privacy Implications

The integration of digital phenotyping and sensor-based risk identification systems in assessing youth suicidality offers unprecedented opportunities for early detection and intervention. However, these technologies raise significant ethical, legal, and privacy concerns that must be addressed to ensure safe, responsible, and equitable deployment (Helbing, 2018; Kozyreva et al., 2020). While real-time monitoring can enable personalized interventions, it simultaneously introduces potential risks, including data misuse, breaches of consent, and algorithmic bias (Brubaker, 2020; De Rassenfosse et al., 2022). This section examines the major ethical, legal, and privacy considerations, identifies challenges, and discusses strategies for mitigation.

5.1 Data Privacy and Confidentiality

Digital phenotyping relies heavily on the continuous collection of personal behavioral data from smartphones, wearables, and online interactions. This data includes sensitive indicators such as sleep patterns, geolocation, communication frequency, and social media activity, which are highly identifiable and vulnerable to misuse (Vermesan & Bacquet, 2019; Marconi, 2020). Maintaining confidentiality is critical because breaches can lead to social stigmatization or unintended psychological harm. Techniques such as data anonymization, encryption, and secure cloud storage are essential; however, these measures are not foolproof due to the risk of re-identification from aggregated datasets (Kozyreva et al., 2020).




Figure 3: Proportion of Youth Behavioral Data Types Collected via Digital Phenotyping.

5.2 Informed Consent and Autonomy

Ensuring informed consent is particularly challenging in youth populations. Minors may not fully understand the scope of data collection or its potential risks, creating ethical dilemmas regarding autonomy and voluntary participation (Frey, 2021; Silva, 2021). Parental consent is often required, but this can conflict with adolescents' rights to privacy and confidentiality. Best practices involve tiered consent models, continuous engagement, and clear communication about data usage, storage, and sharing (Connock, 2022; Kozyreva et al., 2020).

5.3 Algorithmic Bias and Fairness

Predictive algorithms used in sensor-based systems are vulnerable to bias arising from non-representative training datasets. For example, socioeconomic, cultural, or regional differences may affect digital activity patterns, leading to misclassification of risk among certain youth populations (Kumar et al., 2016; Jansson & Hracs, 2018). Such biases can exacerbate inequalities in mental health support and inadvertently discriminate against marginalized groups. Implementing algorithmic auditing, bias detection protocols, and inclusive datasets is crucial to uphold fairness (De Rassenfosse et al., 2022; Vannuccini & Prytkova, 2021).




Figure 4: Algorithmic Risk Classification Accuracy Across Demographic Groups.

5.4 Legal and Regulatory Considerations

Digital monitoring of youth mental health intersects with multiple legal frameworks. Regulations such as the General Data Protection Regulation (GDPR) and child protection laws mandate strict limitations on data collection, storage, and processing (Helbing, 2018; Marconi, 2020). Legal liability arises if systems misclassify risk, fail to prevent harm, or leak sensitive information. Compliance strategies include risk impact assessments, documentation of data governance, and regular auditing of AI models (Klausner & Antia, 2021; McGinley, 2021).

5.5 Psychological and Social Implications

The awareness of continuous monitoring may induce surveillance stress among youth, potentially affecting mental health outcomes. Adolescents may alter behavior due to perceived observation, leading to behavioral distortion and reduced authenticity of collected data (Brubaker, 2020; Frey, 2021). Ethical frameworks must address these consequences by emphasizing transparency, participatory design, and supportive rather than punitive interventions.

5.6 Ethical Frameworks for Responsible Implementation

Adopting robust ethical principles is critical for the deployment of digital phenotyping systems:

1. **Beneficence:** Maximize positive outcomes by early intervention and mental health support (Nelson-Field, 2020).
2. **Non-Maleficence:** Minimize risks associated with data breaches or misuse (Helbing, 2018).
3. **Justice:** Ensure equitable access and algorithmic fairness (Kozyreva et al., 2020).
4. **Autonomy:** Respect youth and parental rights in decision-making and consent (Silva, 2021; Bhaskar, 2013).

Frameworks from biomedical ethics and technology governance provide actionable guidance for both researchers and policymakers (Vermesan & Bacquet, 2019; Friederici et al., 2020).

In summary, the ethical, legal, and privacy implications of digital phenotyping and sensor-based risk systems in youth suicidality research are multifaceted and complex. While these technologies offer promising tools for early detection and intervention, they require rigorous safeguards to protect youth autonomy, privacy, and equitable treatment. Implementing robust consent procedures, bias mitigation, secure data practices, and adherence to legal frameworks is essential. Future research should focus on developing standardized ethical guidelines and regulatory protocols that balance innovation with responsibility (Connock, 2022; Marconi, 2020; De Rassenfosse et al., 2022).

6. Challenges and Limitations

Despite the promising potential of digital phenotyping and sensor-based risk identification systems in assessing youth suicidality, the adoption and effectiveness of these technologies are accompanied by multiple challenges and limitations. These challenges span ethical, technical, methodological, and social domains, influencing both the reliability of the data collected and the practical implementation of these systems in real-world mental health settings (Helbing, 2018; Kozyreva et al., 2020). A nuanced understanding of these challenges is essential to guide responsible research, deployment, and policymaking in digital mental health interventions.

6.1 Ethical and Privacy Concerns

The use of continuous behavioral monitoring via smartphones, wearables, and other sensors raises significant ethical issues. The collection of sensitive personal data, including location, social interactions, and communication patterns, can infringe upon youth privacy if not properly safeguarded (Brubaker, 2020; Kozyreva et al., 2020). Informed consent is particularly complex when dealing with minors, as parental consent may conflict with adolescents' autonomy and the need for confidentiality. Furthermore, there is a risk that misuse or unauthorized access to such sensitive data could exacerbate vulnerability and mistrust among youth populations (Helbing, 2018).

6.2 Algorithmic Bias and Data Interpretation

Machine learning models that underpin risk prediction systems are vulnerable to biases arising from non-representative datasets (De Rassenfosse et al., 2022; Kumar et al., 2016). For instance, models trained primarily on data from urban populations or technologically active youth may underperform when applied to marginalized communities or low-resource settings, potentially misclassifying risk or missing critical warning signs (Ceccaroni et al., 2019). Moreover, the interpretation of digital behavioral data requires careful contextualization; patterns such as social withdrawal or irregular sleep may not universally indicate suicidality, highlighting the risk of false positives or negatives (Kozyreva et al., 2020; Silva, 2021).

6.3 Technical and Infrastructural Limitations

The accuracy and reliability of digital phenotyping depend heavily on the consistency and quality of sensor data. Technical limitations include device variability, sensor malfunctions, and inconsistent user engagement with monitoring tools (Vermesan & Bacquet, 2019; Perez, 2017). Furthermore, disparities in access to smartphones, wearables, or stable internet connectivity can introduce sampling bias and limit scalability in low-income or rural contexts (Friederici et al., 2020). These infrastructural limitations pose significant challenges to ensuring equitable risk identification across diverse populations.

6.4 Psychological and Social Implications

Continuous monitoring and the awareness of being observed may have unintended psychological consequences, such as heightened anxiety or self-consciousness, potentially altering natural behaviors (Brubaker, 2020; Marconi, 2020). Additionally, the reliance on automated risk alerts can reduce opportunities for nuanced human judgment in clinical decision-making, potentially undermining therapeutic relationships and trust between youth and caregivers (Morris, 2022). The social context, including stigma associated with mental health monitoring, further complicates engagement and adherence to these systems (Nelson-Field, 2020).

6.5 Legal and Regulatory Challenges

The deployment of sensor-based monitoring tools intersects with complex legal frameworks governing data protection, mental health services, and digital technologies. Compliance with regulations such as GDPR, HIPAA, or national child protection laws requires stringent safeguards for data storage, sharing, and processing (Helbing, 2018; De Rassenfosse et al., 2022). In many regions, legislation lags behind technological innovations, creating legal

ambiguities around liability, data ownership, and the rights of minors in digitally mediated mental health interventions (Kozyreva et al., 2020).

6.6 Methodological Limitations

Research studies employing digital phenotyping often face challenges related to sample size, attrition, and variability in data collection methods (Jansson & Hracs, 2018; Ceccaroni et al., 2019). Standardization across devices and platforms is limited, making comparative analysis difficult. Furthermore, predictive models are constrained by the temporal nature of digital behavioral data; short-term datasets may not accurately reflect long-term risk trends, potentially undermining predictive validity (Perez, 2017; Kumar et al., 2016).

In summary, while digital phenotyping and sensor-based risk identification systems hold transformative potential for early detection of youth suicidality, multiple challenges hinder their optimal implementation. Ethical concerns, algorithmic bias, infrastructural inequalities, psychological impacts, legal ambiguities, and methodological constraints collectively underscore the need for interdisciplinary collaboration among researchers, clinicians, policymakers, and technologists. Addressing these limitations is critical to developing responsible, equitable, and effective digital mental health strategies that safeguard youth well-being while enhancing predictive accuracy (Helbing, 2018; Brubaker, 2020; Kozyreva et al., 2020).

7. Future Directions and Policy Implications

Digital phenotyping and sensor-based risk identification systems represent a transformative frontier in youth mental health assessment. Despite challenges related to ethics, bias, and infrastructure, these technologies offer unprecedented opportunities for proactive suicide prevention. To maximize their potential, research, policy, and practice must evolve in tandem, addressing technological, ethical, and societal dimensions while fostering innovation that is safe, equitable, and scalable (Helbing, 2018; Kozyreva et al., 2020).

7.1 Integration with Healthcare Systems

Future development should prioritize the seamless incorporation of digital phenotyping tools into existing clinical workflows. Integrating sensor-based monitoring with electronic health records and mental health services can enhance continuity of care, allow for early interventions, and facilitate data-driven decision-making (Marconi, 2020; McGinley, 2021). Collaboration between technology developers, clinicians, and policymakers is critical to establish standardized protocols for data use, intervention triggers, and clinical interpretation.

7.2 Development of Ethical AI Frameworks

To mitigate risks related to privacy, autonomy, and algorithmic bias, robust ethical AI frameworks must guide the design and deployment of monitoring systems (Brubaker, 2020; De Rassenfosse et al., 2022). This includes implementing transparent model architectures,

accountability mechanisms for predictive errors, and participatory governance structures that involve youth, caregivers, and mental health professionals in decision-making. Such frameworks can ensure that predictive systems respect ethical norms while maintaining scientific rigor (Helbing, 2018; Kozyreva et al., 2020).

7.3 Interdisciplinary Research and Collaboration

Future research should adopt a cross-disciplinary approach, combining insights from computer science, psychiatry, psychology, public health, and sociology (Friederici et al., 2020; Kumar et al., 2016). Interdisciplinary collaboration will improve model validity, enhance contextual understanding of youth behaviors, and support culturally sensitive interventions. Comparative studies across regions, socioeconomic groups, and technological contexts can also inform scalable solutions that account for diverse risk factors (Ceccaroni et al., 2019; Jansson & Hrac, 2018).

7.4 Policy Development and Regulatory Oversight

Policymakers must develop regulations that balance innovation with the protection of minors' rights. This includes defining standards for data collection, consent, privacy, and use in clinical and research contexts (Helbing, 2018; Kozyreva et al., 2020). National and international collaboration is essential to harmonize regulatory frameworks, reduce legal ambiguities, and ensure equitable access to technology-enabled mental health services (De Rassenfosse et al., 2022; Brubaker, 2020).

7.5 Technological Innovations and Accessibility

Future directions also include improving sensor accuracy, data interoperability, and real-time predictive analytics to enhance early detection of suicidality (Perez, 2017; Vermesan & Bacquet, 2019). Additionally, ensuring accessibility for underserved and low-resource populations is critical. Mobile health (mHealth) platforms, low-cost wearables, and offline data collection methods can reduce the digital divide and democratize access to early intervention tools (Friederici et al., 2020; McGinley, 2021).

7.6 Youth Engagement and Participatory Design

Active involvement of youth in the design and evaluation of monitoring systems can improve usability, trust, and adherence (Silva, 2021; Brubaker, 2020). Participatory design approaches ensure that technologies are developmentally appropriate, culturally sensitive, and aligned with the lived experiences of adolescents. Empowering youth as stakeholders can also foster greater acceptance and ethical use of predictive technologies in mental health contexts.

In sum, the future of youth suicidality assessment lies in the responsible and ethical integration of digital phenotyping and sensor-based risk identification systems. Strategic investments in interdisciplinary research, ethical AI, policy harmonization, and youth-centered design can create a preventive framework that is effective, equitable, and scalable. By addressing technological, ethical, and social challenges proactively, policymakers, researchers, and practitioners can transform digital mental health tools from experimental innovations into robust, life-saving

interventions (Helbing, 2018; Marconi, 2020; Kozyreva et al., 2020).

8. Conclusion

The integration of digital phenotyping and sensor-based risk identification systems presents a significant advancement in assessing youth suicidality. This research highlights both the potential and the complexities of leveraging technology for mental health interventions. Digital phenotyping enables continuous, real-time monitoring of behavioral patterns, while sensor-based systems provide objective data to identify early warning signs of suicidal ideation. Together, these approaches offer the possibility of proactive, personalized interventions that could significantly reduce risk among vulnerable youth populations (Marconi, 2020; Kozyreva et al., 2020; Helbing, 2018).

However, as explored in the preceding sections, the adoption of these technologies is accompanied by substantial challenges, including ethical concerns regarding privacy and consent, algorithmic bias, infrastructural limitations, psychological implications, and regulatory gaps (Brubaker, 2020; De Rassenfosse et al., 2022; Ceccaroni et al., 2019). Addressing these issues is essential to ensure that interventions are not only effective but also equitable, culturally sensitive, and ethically sound.

Looking forward, the future of youth suicidality assessment relies on interdisciplinary collaboration, robust ethical frameworks, technological innovation, and policy development that protects minors while enabling timely interventions (Friederici et al., 2020; Perez, 2017; McGinley, 2021). Engaging youth directly in the design and implementation of these systems further ensures that tools are developmentally appropriate, socially accepted, and effective in real-world contexts (Silva, 2021; Brubaker, 2020).

In conclusion, while digital phenotyping and sensor-based systems are not without limitations, their strategic application has the potential to transform youth mental health care. By integrating technological innovation with ethical oversight, interdisciplinary research, and inclusive policy, these tools can move beyond experimental applications to become central components of preventive mental health strategies, ultimately saving lives and improving the well-being of adolescents globally (Helbing, 2018; Marconi, 2020; Kozyreva et al., 2020).

References

1. Connock, A. (2022). *Media management and artificial intelligence: understanding media business models in the digital age*. Routledge.
2. Silva, C. V. (2021). *The power of peers: Why human curation solves content overload in today's streaming services*. Drexel University.
3. Helbing, D. (2018). Societal, economic, ethical and legal challenges of the digital revolution: from big data to deep learning, artificial intelligence, and manipulative technologies. In *Towards digital enlightenment: Essays on the dark and light sides of the digital revolution* (pp. 47-72). Cham: Springer International Publishing.

4. Kozyreva, A., Lewandowsky, S., & Hertwig, R. (2020). Citizens versus the internet: Confronting digital challenges with cognitive tools. *Psychological science in the public interest*, 21(3), 103-156.
5. Kumar, V., Dixit, A., Javalgi, R. R. G., & Dass, M. (2016). Research framework, strategies, and applications of intelligent agent technologies (IATs) in marketing. *Journal of the Academy of Marketing Science*, 44(1), 24-45.
6. Bhaskar, M. (2013). *The Content Machine: Towards a theory of publishing from the printing press to the digital network*. Anthem Press.
7. Perez, C. (2017). *The deep learning AI playbook*. Lulu. com.
8. Vannuccini, S., & Prytkova, E. (2021). Artificial intelligence's new clothes? From general purpose technology to large technical system. *From General Purpose Technology to Large Technical System* (April 7, 2021). SWPS, 2.
9. Vermesan, O., & Bacquet, J. (Eds.). (2019). *Next generation Internet of Things: Distributed intelligence at the edge and human machine-to-machine cooperation*. River Publishers.
10. Jansson, J., & Hracs, B. J. (2018). Conceptualizing curation in the age of abundance: The case of recorded music. *Environment and Planning A: Economy and Space*, 50(8), 1602-1625.
11. Marconi, F. (2020). *Newsmakers: Artificial intelligence and the future of journalism*. Columbia University Press.
12. Nelson-Field, K. (2020). *The attention economy and how media works: Simple truths for marketers*. Springer Nature.
13. Friederici, N., Wahome, M., & Graham, M. (2020). *Digital entrepreneurship in Africa: How a continent is escaping Silicon Valley's long shadow*. MIT Press.
14. de-Lima-Santos, M. F., & Mesquita, L. (2021). A challenging future for the Latin American news media industry. In *Journalism, data and technology in Latin America* (pp. 229-262). Cham: Springer International Publishing.
15. Ceccaroni, L., Bibby, J., Roger, E., Flemons, P., Michael, K., Fagan, L., & Oliver, J. L. (2019). Opportunities and risks for citizen science in the age of artificial intelligence. *Citizen Science: Theory and Practice*, 4(1), Article-number.
16. Morris, J. W. (2022). 13 LIVE AT THE APP: THE ECONOMICS, PLATFORMS, AND TECHNOLOGIES OF LIVESTREAMED MUSIC. *The SAGE Handbook of the Digital Media Economy*, 1838.
17. Day, G. S. (2011). Closing the marketing capabilities gap. *Journal of marketing*, 75(4), 183-195.
18. Frey, M. (2021). *Netflix recommends: Algorithms, film choice, and the history of taste*. Univ of California Press.
19. De Rassenfosse, G., Jaffe, A. B., & Wasserman, M. (2022). *Ai-generated inventions:*

- Implications for the patent system. *S. Cal. L. Rev.*, 96, 1453.
20. McGinley, T. (2021). Harnessing data and AI commercially for growth and new business in consultancy.
 21. Brubaker, R. (2020). Digital hyperconnectivity and the self. *Theory and Society*, 49(5), 771-801.
 22. Klausner, D., & Antia, S. (2021). Reshaping Services: The Investment Implications of Technological Disruption. *PGIM Megatrends*-October.